

## Inflammatory response to assess toxicity of biodiesel emission samples

Christoph Vogel, PhD  
Environmental Toxicology  
Center for Health and the Environment  
University of California Davis

ARB December 8, 2010  
Biodiesel and Renewable Diesel Multimedia  
Evaluation Public Meeting

---

---

---

---

---

---

---

## Inhalation of environmental and occupational pollutants in vivo

- Pulmonary inflammation, fibrosis, lung carcinomas
- Cardiovascular diseases like atherosclerosis
- Chronic inflammatory response as a main cause for adverse health effects

---

---

---

---



---

---

---

## In vitro cell models

- Two main target cell types
  - a) Macrophages (U937), phagocyte, acts as first line of defense
  - b) Lung Clara cells from pulmonary epithelium (NCI H441)



---

---

---

---

---

---

---

## Biomarkers of PM exposure, inflammation and oxidative stress

- CYP1A1: Cytochrome P450 monooxygenase, xenobiotic metabolizing enzyme, Arylhydrocarbon-Receptor regulated
- COX-2: Cyclooxygenase, key enzyme for production of prostaglandins involved in inflammation
- IL-8: Interleukin 8, chemoattractant peptide for neutrophils, major mediator of inflammatory response
- HO-1: Hemeoxygenase 1, essential enzyme in heme catabolism, protect cells against oxidative injury. Induced by exposure to various forms of oxidative stress

---

---

---

---


---

---

---

---

## Macrophage Model to measure Inflammation caused by Diesel PM




---

---

---

---


---

---

---

---

## Formation of foam cells




---

---

---

---


---

---

---

---

## Development of atherosclerotic lesions in ApoE mice




---

---

---

---

---


---

---

---

## Long-term Air Pollution Exposure and Acceleration of Atherosclerosis and Vascular Inflammation in an Animal Model

6 hrs/day, 5 days/week x 6 months  
Mean levels only 15.2  $\mu\text{g}/\text{m}^3$



Air pollution exposed mice developed more ATHEROSCLEROSIS

JAMA. 2005;294:3003-3010

---

---

---

---


---

---

---

---

## Lung Clara cell model (NCIH441)



- Chronic obstructive pulmonary disease (COPD)
- Emphysema
- Asthma

---

---

---

---


---

---

---

---

### Draft C15 Emission Rate: Macrophage CYP1a1 response per mile




---

---

---

---


---

---

---

---

### Draft C15 Emission Rate: Lung Clara cells CYP1a1 response per mile




---

---

---

---


---

---

---

---

### Draft C15 Emission Rate: Lung Clara cells COX-2 response per mile




---

---

---

---

---


---

---

---



### Draft C15 Emission Rate: Macrophage IL-8 response per mile




---

---

---

---


---

---

---

---

### Draft C15 Emission Rate: Lung Clara cells MUC5AC response per mile




---

---

---

---


---

---

---

---

### Draft C15 Emission Rate: Macrophage HO-1




---

---

---

---

---

---

---

---

## Comet Assay or Single-Cell-Gel-Electrophoresis assay

- sensitive technique for the detection of DNA damage at the level of the individual eukaryotic cell
- standard technique for evaluation of DNA damage, biomonitoring genotoxicity

---

---


---

---

---

---

---



---

---

---

---

---

---

---

## Comet Standard Cells 1



---

---

---


---

---

---

---

Comet Standard Cells 2



---

---

---

---


---

---

---

---

Comet Standard Cells 3



---

---

---

---


---

---


---

---

Control



Carb



Undamaged DNA retains a highly organized association with matrix proteins in the nucleus

---

---

---

---


---

---


---

---

Soy 100



Animal 100



---

---

---


---

---


---

---

Renewable 100



NIST SRM1650



Circular "head" corresponding to the undamaged DNA and a "tail" of damaged DNA,  
the brighter and longer the tail, the higher the level of damage

---

---

---


---

---

---

---

NIST SRM1650



---

---

---


---

---

---

---

## DNA damage measured by the comet assay



Percent Tail DNA was measured after 3-h treatment of U937 cells under serum-free conditions with 200 µg/ml extracts of PMs.

## Summary

- Carb and Biodiesel blends induce CYP1A1 through PAHs which bind to and activate the Ah-Receptor
- Carb and Biodiesel blends induce inflammatory markers like COX-2 and IL-8 in macrophages and MUC5AC in lung Clara cell type (NCI H441)
- Effect of Biodiesel blends on inflammatory markers like COX-2 and IL-8 tend to be lower than Carb diesel
- No genotoxic effects of biodiesel blends in Comet assay

## Thank you

Bob Okamoto  
 Norm Kado  
 Reiko Kobayashi  
 Xiaoxue Liu  
 Dalei Wu  
 Wen Li  
 Viktoria Kuo  
 Patty Lok  
 Danitza Alvizar  
 Helen Woldai  
 Pat Wong  
 Fumio Matsumura